

# ODD-GRACEFUL LABELINGS OF TREES OF DIAMETER 5

CHRISTIAN BARRIENTOS

**ABSTRACT.** A difference vertex labeling of a graph  $G$  is an assignment  $f$  of labels to the vertices of  $G$  that induces for each edge  $xy$  the weight  $|f(x) - f(y)|$ . A difference vertex labeling  $f$  of a graph  $G$  of size  $n$  is odd-graceful if  $f$  is an injection from  $V(G)$  to  $\{0, 1, \dots, 2n - 1\}$  such that the induced weights are  $\{1, 3, \dots, 2n - 1\}$ . We show here that any forest whose components are caterpillars is odd-graceful. We also show that every tree of diameter up to five is odd-graceful.

## 1. INTRODUCTION

Let  $G$  be a graph of order  $m$  and size  $n$ , a *difference vertex labeling* of  $G$  is an assignment  $f$  of labels to the vertices of  $G$  that induces for each edge  $xy$  a label or *weight* given by the absolute value of the difference of its vertex labels. Graceful labelings are a well-known type of difference vertex labeling; a function  $f$  is a *graceful labeling* of a graph  $G$  of size  $n$  if  $f$  is an injection from  $V(G)$  to the set  $\{0, 1, \dots, n\}$  such that, when each edge  $xy$  of  $G$  has assigned the weight  $|f(x) - f(y)|$ , the resulting weights are distinct; in other words, the set of weights is  $\{1, 2, \dots, n\}$ . A graph that admits a graceful labeling is said to be *graceful*.

When a graceful labeling  $f$  of a graph  $G$  has the property that there exists an integer  $\lambda$  such that for each edge  $xy$  of  $G$  either  $f(x) \leq \lambda < f(y)$  or  $f(y) \leq \lambda < f(x)$ ,  $f$  is named an  $\alpha$ -labeling and  $G$  is said to be an  $\alpha$ -graph. From the definition it is possible to deduce that an  $\alpha$ -graph is necessarily bipartite and that the number  $\lambda$  (called the *boundary value* of  $f$ ) is the smaller of the two vertex labels that yield the edge with weight 1. Some examples of  $\alpha$ -graphs are the cycle  $C_n$  when  $n \equiv 0 \pmod{4}$ , the complete bipartite graph  $K_{m,n}$ , and caterpillars (i.e., any tree with the property that the removal of its end vertices leaves a path).


A little less restrictive than  $\alpha$ -labelings are the odd-graceful labelings introduced by Gnanajothi in 1991 [4]. A graph  $G$  of size  $n$  is *odd-graceful* if there is an injection  $f : V(G) \rightarrow \{0, 1, 2, \dots, 2n - 1\}$  such that the set of induced weights is  $\{1, 3, \dots, 2n - 1\}$ . In this case,  $f$  is said to be an *odd-graceful labeling* of  $G$ . One of the applications of these labelings is that trees of size  $n$ , with a suitable odd-graceful labeling, can be used to generate cyclic decompositions of the complete bipartite graph  $K_{n,n}$ . In Figure 1 we show an odd-graceful tree of size 6 together with its embedding in the circular arrangement used to produce the cyclic decomposition of  $K_{6,6}$ . Once the labeled tree has been embedded, successive  $30^\circ$  (counterclockwise) rotations produce the desired cyclic decomposition of  $K_{6,6}$ .

---

2000 *Mathematics Subject Classification.* Primary 05C78.

*Key words and phrases.* Odd-graceful labeling,  $\alpha$ -labeling, trees of diameter 5.

This paper is in final form and no version of it will be submitted for publication elsewhere.

Fig. 1. Cyclic decomposition of  $K_{6,6}$ 

Gnanajothi [4] proved that the class of odd-graceful graphs lies between the class of  $\alpha$ -graphs and the class of bipartite graphs; she proved that every  $\alpha$ -graph is also odd-graceful. The reverse case does not work, for example the odd-graceful tree shown in Figure 1 is the smallest tree without an  $\alpha$ -labeling. Since many families of  $\alpha$ -graphs are known, the most attractive examples of odd-graceful graphs are those without an  $\alpha$ -labeling or where an  $\alpha$ -labeling is unknown; for instance, Gnanajothi [4] proved that the following are odd-graceful graphs:  $C_n$  when  $n \equiv 2(\text{mod } 4)$ , the disjoint union of  $C_4$ , the prism  $C_n \times K_2$  if and only if  $n$  is even, and trees of diameter 4 among others. Eldergil [2] proved that the one-point union of any number of copies of  $C_6$  is odd-graceful. Seoud, Diab, and Elsakhawi [5] showed that a connected  $n$ -partite graph is odd-graceful if and only if  $n = 2$  and that the join of any two connected graphs is not odd-graceful.

A detailed account of results in the subject of graph labelings can be found in Gallian's survey [3].

Gnanajothi [4] conjectured that all trees are odd-graceful and verified this conjecture for all trees with order up to 10. The author has extended this up to trees with order up to  $12^1$ . In this paper we prove that all trees of diameter 5 are odd-graceful and that any forest whose components are caterpillars is odd-graceful.

## 2. ODD-GRACEFUL FORESTS

In this section we study *forests* that accept odd-graceful labelings. Recall that a forest with more than one component cannot be graceful because it has "too many edges". First we prove that any graph that admits an  $\alpha$ -labeling also admits an odd-graceful labeling by transforming conveniently its  $\alpha$ -labeling.

**Theorem 1.** *Any  $\alpha$ -graph is odd-graceful.*

*Proof.* Let  $G$  be an  $\alpha$ -graph of size  $n$ , as consequence  $G$  is bipartite with partition  $\{A, B\}$ . Suppose that  $f$  is an  $\alpha$ -labeling of  $G$  such that  $\max\{f(x) : x \in A\} < \min\{f(x) : x \in B\}$ . Let  $g$  be a labeling of the vertices of  $G$  defined by

$$g(x) = \begin{cases} 2f(x), & x \in A \\ 2f(x) - 1, & x \in B. \end{cases}$$

Thus, the labels assigned by  $g$  are in the set  $\{0, 1, \dots, 2n - 1\}$ , furthermore, the weight of the edge  $xy$  of  $G$  induced by the labeling  $f$ , where  $x \in A$  and  $y \in B$ , is  $w = f(y) - f(x)$ , so its weight under the labeling  $g$  is  $g(y) - g(x) = 2f(y) - 1 -$

<sup>1</sup>Odd-graceful labelings of trees of order 11 and 12 can be found at <http://cims.clayton.edu/cbarrien/research>

$2f(x) = 2(f(y) - f(x)) - 1 = 2w - 1$ . Since  $1 \leq w \leq n$ , we have that the weights induced by  $g$  are  $\{1, 3, \dots, 2n - 1\}$ . Therefore,  $g$  is an odd-graceful labeling of  $G$ .  $\square$

In Figure 2 we show an example of an  $\alpha$ -labeling of a caterpillar, followed for the corresponding odd-graceful labeling. We use this labeling in the next theorem.




Fig. 2. Odd-graceful labeling of a caterpillar

**Theorem 2.** *Any forest which components are caterpillars is odd-graceful.*

*Proof.* Let  $F_i$  be a caterpillar of size  $n_i \geq 1$ , for  $1 \leq i \leq k$ . Let  $u_i, v_i \in V(F_i)$  such that  $d(u_i, v_i) = \text{diam}(F_i)$ ; so identifying  $v_i$  with  $u_{i+1}$ , for each  $1 \leq i \leq k - 1$ , we have a caterpillar  $F$  of size  $\sum_{i=1}^k n_i = n$ . Now we proceed to find both, the  $\alpha$ -labeling of  $F$  and its corresponding odd-graceful labeling, using the scheme shown in Figure 2. Once the odd-graceful labeling has been obtained, we disengage each caterpillar  $F_i$  from  $F$ , keeping their labels; in this form, the weights induced are  $\{1, 3, \dots, 2n - 1\}$ . To eliminate the overlapping of labels we subtract 1 from each vertex label of  $F_i$  when  $i$  is even, in this way the weights remain the same and the labels assigned on  $u_{i+1}$  and  $v_i$  differ by one unit. Therefore, the labeling of the forest  $\bigcup_{i=1}^k F_i$  is odd-graceful.  $\square$

In Figure 3 we show an example of this construction using the odd-graceful labeling obtained in Figure 2.




Fig. 3. Odd-graceful labeling of a caterpillar

The procedure used in this proof can be extended to the disjoint union of graphs with  $\alpha$ -labelings. In fact, suppose that the concatenation of blocks  $B_1, B_2, \dots, B_k$  results in a graph  $G$  whose block-cutpoint graph is a path. In [1] we proved that if each  $B_i$  is an  $\alpha$ -graph, so it is  $G$ . Transforming this  $\alpha$ -labeling into an odd-graceful labeling and disconnecting  $G$  into blocks, the disjoint union of these blocks is odd-graceful.

**Theorem 3.** *The disjoint union of blocks that accept  $\alpha$ -labelings is odd-graceful.*

### 3. ODD-GRACEFUL TREES OF DIAMETER FIVE

Every tree of diameter at most 3 is a caterpillar, therefore it is odd-graceful. Gnanajothi [4] proved that every rooted tree of height 2 (that is, diameter 4) is odd-graceful. In the next theorem we represent trees of diameter 5 as rooted trees of height 3 and prove that they are odd-graceful.

Let  $T$  be a tree of diameter 5;  $T$  can be represented as a rooted tree of height 3 by using any of its two central vertices as the root vertex. Note that only one of the vertices in level 1 has descendants in level 3; this vertex will be located in the right extreme of level 1. Now, within each level, the vertices are placed from left to right in such a way that their degrees are increasing. In the proof of the next theorem we use this type of representation of  $T$ , that is, assuming that  $v$  (one of the two central vertices) is the root.

**Theorem 4.** *All trees of diameter five are odd-graceful.*

*Proof.* Let  $T$  be a tree of diameter 5 and size  $n$ . Suppose that  $T$  has been drawn according to the previous description. Let  $v_{i,j}$  denote the  $i$ th vertex of level  $j$ , for  $j = 1, 2, 3$ , this vertex is placed at the right of  $v_{i+1,j}$ . Consider the labeling  $f$  of the vertices within each level given by recurrence as follows:  $f(v) = 0$ ,  $f(v_{1,1}) = 2n - 2\deg(v) + 1$ ,  $f(v_{1,2}) = 2$ ,  $f(v_{1,3}) = 3$ , and  $f(v_{i,j}) = f(v_{i-1,j}) + d(v_{i,j}, v_{i-1,j})$  where  $i \geq 2$  and  $1 \leq j \leq 3$ .

We claim that  $f$  is an odd-graceful labeling of  $T$ . In fact, let us see that there is no overlapping of labels. On level 0 the label used is 0 and on level 2 all labels are even being 2 the smallest label used here. On levels 1 and 3 the labels used are odd; on level 1 the labels used are  $2n - 1, 2n - 3, \dots, 2n - 2\deg(v) + 1$ , while on level 3 the labels used are  $3, 5, \dots, 2\deg(v_{1,2}) - 1$ . Now we need to prove that  $2n - 2\deg(v) + 1 > 2\deg(v_{1,2}) - 1$ ; since  $T$  is a tree of diameter 5, at least two vertices on level 1 has descendants, so  $n + 1 > \deg(v) + \deg(v_{1,2})$ , which implies the desired inequality.

As a consequence of the fact that labels used in consecutive levels have different parity, each weight obtained is an odd number not exceeding  $2n - 1$ . Suppose that  $v_{i+1,j}$  and  $v_{i,j}$  have the same father  $x$ , by definition of  $f$ , the edges  $xv_{i+1,j}$  and  $xv_{i,j}$  have consecutive weights. If  $v_{i+1,j}$  and  $v_{i,j}$  have different father,  $x$  and  $y$  respectively, then  $|f(y) - f(v_{i,j})| = |(f(x) + 2) - (f(v_{i+1,j}) + 4)| = |f(x) - f(v_{i+1,j}) - 2|$ . Thus, the weights are  $2n - 2\deg(v) - 1, \dots, 2\deg(v) + 1$ . On level 2, the weights are  $2n - 2\deg(v) - 3, \dots, 2\deg(v_{1,2}) - 1$ , and on level 3 the weights are  $2\deg(v_{1,2}) - 3, \dots, 1$ .

Therefore,  $f$  is an odd-graceful labeling of  $T$ . □

In Figure 4 we present a scheme of this labeling for a tree of size 13.




Fig. 4. Odd-graceful tree of diameter 5

Similar arguments can be used to find odd-graceful labelings of trees of diameter 6; however we do not have a general labeling scheme for this case. So it is an open problem determining whether trees of diameter 6 are odd-graceful. In Figure 5, we give an example of an odd-graceful labeling for a tree of size 17 and diameter 6.




Fig. 5. Odd-graceful tree of diameter 6

To conclude this section, we show in Figure 6 an odd-graceful labeling for a special type of tree of diameter 6, namely the star  $S(n, 3)$  with  $n$  spokes of length 3.




Fig. 6. Odd-graceful labeling of the star  $S(n, 3)$

## REFERENCES

- [1] C. Barrientos, Graceful labelings of chain and corona graphs, *Bull. Inst. Combin. Appl.*, **34**(2002) 17-26
- [2] P. Eldergill, Decomposition of the Complete Graph with an Even Number of Vertices. M. Sc. Thesis, McMaster University, 1997
- [3] J.A. Gallian, A dynamic survey of graph labeling, *Electron. J. Combinatorics* (2008), #DS6
- [4] R.B. Gnanajothi, Topics in Graph Theory. Ph. D. Thesis. Madurai Kamaraj University, 1991
- [5] M.A. Seoud, A.E.I. Abdel Maqsooud, and E.A. Elsayhawi, On strongly- $C$  harmonious, relatively prime, odd graceful and cordial graphs, *Proc. Math. Phys. Soc. Egypt*, **73**(1998) 33-55

DEPARTMENT OF MATHEMATICS, CLAYTON STATE UNIVERSITY, MORROW, GA 30260, USA

*E-mail address:* chr\_barrientos@yahoo.com

*URL:* <http://cims.clayton.edu/cbarrien>